



Fatty acid composition of the marine micro alga *Tetraselmis chuii* Butcher in response to culture conditions

Mehdi Mohammadi^{1,2*}, Najmeh Kazeroni³, Mehran Javaheri Baboli³

1. Department of Marine Biotechnology and Environment, Persian Gulf Research and studies center, Persian Gulf University, Bushehr, Iran,

2-Algae Research Development Co., Persian Gulf St. Bushehr, Iran

3-Department of Fisheries, Science and Research khuzestan Branch, Islamic Azad University, Ahvaz, Iran, E-

3- Department of fisheries science, Ahvaz branch, Islamic Azad University. Khuzestan, Iran,

*mmohammadi@pgu.ac.ir

Abstract

Micro alga *Tetraselmis chuii* Butcher is one of the species that are considered to be important sources of polyunsaturated fatty acids (PUFA). The aim of our experiment was to determine the effect of the light intensity and the water salinity on the growth and fatty acid composition of *Tetraselmis chuii*. There were significant correlation ($P \leq 0.05$) between the cell density and light intensity and cell grow rate and light intensity, the correlation were $r^2=0.71$; $r^2=0.74$ correspondingly. The lowest concentration of the cell density was observed at the lowest light intensity and the highest salinity. Palmitic acid among the saturated fatty acid, oleic acid, among mono unsaturated fatty acid, alpha-linoleic acid among the PUFA, and eicosapentaenoic acid (EPA) among high unsaturated fatty acid (HUFA), had the highest percentage. Variance analyses showed that the light intensity exhibited a significant correlation ($P \leq 0.05$) with PUFA, HUFA. However there was not significant correlation among light intensity and SFA, MUFA. This result showed that relationship between environment conditions and fatty acid composition in *T. chuii* was significant thus it can be used to produce specific profile of fatty acids in this strain.

Keywords : *Tetraselmis chuii*, cell density, light intensity, salinity and fatty acid.

Introduction

Microalgae are the major food source for many aquatic organisms and the main live feed component in marine hatchery operations because they serve as a natural resource for polyunsaturated fatty acids (Makridis *et al.*,2006). *Tetraselmis chuii* Butcher is a green four-flagellated prasinophyte characterized by an ovoid body shape and a distinct curved body when viewed side ways. The micro alga measures 12-14 μ m in length,9-10 μ m in width and belongs to the family Chlamydomonadaceae. *Tetraselmis* is a sizeable genus (more than 50 species) of green flagellates. Most species are known from inshore marine environments, tide pools in particular, but a few freshwater species are also known (Bold and Wynne,1985). The pyrenoid is embedded in the single chloroplast, which occupies most of the volume of the cell, especially near its proximal end (Smith,1955). *Tetraselmis sp* are ideal for culture because they are euryhaline and eurythermal (Fabregas *et al.*,1984).Environmental factors such as temperature, salinity, pH, and light have been reported to affect microalgae growth (Creswell,2010). Light conditions are the main factors affecting microalgae physiology and the most important factor affecting microalgae photosynthesis kinetics. Such changes in the salinity of water often affect the growth, metabolism and photosynthesis of phytoplanktons (Moisander *et al.*,2002;Lartigue *et al.*,2003). Salt might have a direct effect upon processes involved in electron transport and / or photophosphorylation and result in a decreased in the quantum efficiency of photosynthesis (Seeman and Critchley,1985).Whereas high light intensity decreases total polar lipid content with a concomitant increase in the amount of neutral storage lipids, mainly TAG,s (Brown *et al.*,1996). Neutral lipids, such as TG, are storage substance and energy sources, While polar lipid, glycolipids and phospholipids are the structural components of cellullar membranes as well as modulators of photosystem efficiency and regulators of energy flow(Thompson,1996).*Tetraselmis chuii* is one of the species of microalgae that is most extensively used in aquaculture and is considered to be an optimal source of long-chain PUFAs, and especially of eicosapentaenoic acid (EPA) (Meseck *et al.*,2005;Zaki and Saad,2010).The present study focuses on the adaptation of *Tetraselmis chuii* to varied range of salinity and different light intensity conditions and their effect on the growth and fatty acid composition.

Materials and Methods

Growth conditions

This study provides for to determine effect different light intensities, 2500,4500,6500 lux and the degree of the water salinity 20,25,35,40 ppt on the growth and fatty acid composition of *Tetraselmis chuii* in Conway medium. All the samples were incubated at constant temperature of 25-27°C and pH of 7.5 to 8.2, creating a growth atmosphere that encouraged the growth of the sample daily. Light intensity was measured using a photometer (Model Lutran lx-107). Cell counts of *T.chuii* were measured using a hemocytometer. The specific growth rates (μ) of samples were determined with the following exponential growth equation: (Guillard, 1973).

$$\mu = \ln(N(t) - N_0) / (t - t_0)$$

Table 1. Chemical compounds utilized in Conway medium preparation.

Chemical compound	Concentration
NaNO ₃	116 g
Na ₂ EDTA	45 g
H ₃ Bo3	33.6 g
Na ₂ H ₂ Po ₄ .4H ₂ O	20 g
Fecl ₃ .6H ₂ O	1.3 g
Mncl ₂ .4H ₂ o	0.36 g
Zncl ₂	2.1 g
CoCl ₂ .6H ₂ o	2 g
(NH ₄) ₆ MoO ₇ .4H ₂ O	0.9 g
Cuso ₄ .5H ₂ o	2 g
Vitamin B1	200 mg
Vitamin B12	10 mg


After reaching the end of an exponential phase growth, 800 ml of sample was harvested and centrifuged (model 5810R appendrof) at 2500 rpm, for 5 min, Prior to analysis they were frozen at -70 °c.

Lipid extraction


Total lipids were extracted from 1g of moist sample according to (Folch *et al.*,1957). All samples were dissolved in chloroform and methanol in ratio of 2:1. Chloroform, methanol were allowed to evaporate under nitrogen gas. The fatty acid methyl esters were prepared by direct esterification (Desvilettes *et al.*,1994) of lipid extracts. The lipid extracts were saponified in methanolic KOH (2N) for 10 min at 60°C; conversion into fatty acid methyl esters (FAME) was performed by using methanolic H₂SO₄. Separation and identification of the component fatty acids were done using a GC-Agilent-6890, as liquid chromatograph with a capillary column (30 m length, 0.25 mm i.d , 0.25 μ m film thickness) using nitrogen as carrier gas at 1.45ml/min. The collected data were analyzed using one-way analysis of variance (ANOVA). Significant differences among the different treatments were determined using the Tukey multiple range test at 0.05 level of probability.

Results


Data in table 2 shows the effect salinity and light intensity different on growth of *T.chuii*. The tests were carried out in eighth day when the alga reached maximum cell density (Figure 1). Changes in light quantity and salinity bring about the differences in fatty acids composition (Table 4). The most of SFA were C16:0, which accounted for the highest rate (28.65%) in the sample 25 ppt, 6500 lux, and the lowest rate (19.17%) in the sample 20ppt, 2500 lux determined. Ratio of PUFA to SFA was more in the sample 4500 lux, 25 ppt-6500 lux, 40ppt. The high amount of C20:4n-6 was obtained in 4500 lux,25 ppt. Among mono unsaturated fatty acids, C18:1n-9 was the most dominating. Highest percentages of n-3 showed in 6500 lux, 40 ppt (range 24.81%) and highest percentage of n-6 in 4500 lux,20ppt (range 22.50%).



a



b



c

Figure 1: Changes in cell density of *T.chuii* grown at three different light intensity and four different salinity : 2500lux,20,25,35,40ppt (a), 4500lux,20,25,35,40ppt (b), 6500lux,20,25,35,40ppt (c)

Table 2: Cell density ($\times 10^4$ cell ml⁻¹) and growth rate (division (div) day⁻¹) of *Tetraselmis chuii* exposed to different salinity and light in the Exponential phase.

Light intensity	Salinity	Cell density	Growth rate
2500	20	210 ± 23.3	0.40±0.08
2500	25	262 ± 13	0.44±0.06
2500	35	247 ± 16.8	0.52±0.01
2500	40	206 ± 9.3	0.49±0.03
4500	20	258 ± 27.2	0.48±0.04
4500	25	260 ± 29.4	0.49±0.04
4500	35	261 ± 28.4	0.55±0.04
4500	40	199 ± 25.1	0.51±0.04
6500	20	314 ± 20.3	0.55±0.07
6500	25	324 ± 35.3	0.55±0.04
6500	35	280 ± 29.8	0.55±0.02
6500	40	268 ± 5.1	0.54±0.02

mean ±Standard deviations(n=3).

Table3: Result of one way analysis of variance ANOVA.

Sample	Light	Salinity
	4	y
Cell density		0.05*
Total fatty acid	0.07	0.7
∑ SFA	0.8	0.8
∑ MUFA	0.01*	0.9
∑ PUFA	0.01*	0.7
∑ HUFA	0.04*	0.7
∑ (n-3)	0.04*	0.9
∑ (n-6)	0.04*	0.8

*Significantly different at $P \leq 0.05$ indicate.

Table4: Fatty acids composition (percent of total fatty acids of fresh weight) of microalgae *T.chuii* at the exponential phase

Sample	2500lux 2500lux 20ppt	2500lux 25ppt	2500lux 35ppt	2500lux 40ppt	4500lux 20ppt	4500lux 25ppt	4500lux 35ppt	4500lux 40ppt	6500lux 20ppt	6500lux 25ppt	6500lux 35ppt	6500lux 40ppt
Saturated												
C14:0	0.72	0.69	1.03	0.75	1.23	0.44	0.44	1.65	0.91	0.89	0.95	0.56
C16:0	19.17	26.84	28.81	25.95	24.36	23.51	22.59	27.75	27.48	28.65	27.06	23.41
C18:0	0.72	0.71	0.51	0.98	1.28	0.5	0.82	3.68	0.75	0.79	0.29	0.51
C20:0	3.83	2.68	2.68	3.41	2.55	2.61	2.67	1.51	3.57	4.82	4.21	0.26
C22:0	0.07	0.13	0.09	0.07	0.22	0.11	2.06	1.06	0.08	0.34	0.11	0.03
Monosaturated												
C14:1n5	0.15	0.09	0.15	0.08	0.04	0.01	0.03	0.25	0.15	0.16	0.15	0.11
C16:1n7	3.72	3.79	4.12	2.63	1.4	1.33	2.86	4.1	1.23	1.73	1.73	0.89
C18:1n9	31.37	25.52	21.32	21.61	11.41	12.13	24.33	14.98	13.61	15.38	13.31	14.63
Polyunsaturated												
C18:2n6	7.37	6.54	9.11	8.68	18.52	16.79	8.61	13.48	10.45	8.79	10.24	10.33
C18:3n3	12.86	10.36	7.43	10.71	11.41	10.92	13.61	6.43	12.68	11.39	14.26	12.43
C18:3n6	0.1	0.24	0.16	0.06	0.22	0.18	0.05	0.27	0.21	0.26	0.26	0.04
C18:4n3	0.08	0.3	0.22	0.14	1.2	1.55	0.04	1.08	0.09	1.86	0.05	0.04
C20: 3n6	1.1	1.93	1.34	1.77	0.36	0.14	0.07	0.22	1.62	1.77	1.51	2.7
C20:3n3	0.96	1.01	1.48	0.91	0.06	0.05	0.01	0.06	0.05	0.24	0.08	0.21
C20:4n6	0.36	0.23	0.2	0.12	3.34	4.74	1.69	1.18	1.95	0.2	1.49	2.83
Highunsaturated												
C20:5n3	3.66	4.37	4.52	3.72	4.52	6.56	5.34	3.05	5.99	6.55	6.83	12.22
C22:5n6	0.13	0.23	0.01	0.12	0.04	0.06	0.04	0.26	0.07	0.04	0.04	0.06
C22:5n3	0.26	0.03	0.02	0.08	0.03	0.04	0.03	0.05	0.05	0.09	0.02	0.02
C22:6n3	0.18	0.14	0.11	0.08	0.12	0.19	0.12	0.3	0.08	0.2	0.06	0.13
Sum	86.88	85.92	83.4	81.95	82.4	81.95	85.52	81.54	81.1	84.22	82.74	81.48
∑SFA	24.52	31.07	33.05	31.17	29.66	27.69	28.60	35.67	32.79	35.51	32.64	24.79
∑MUFA	35.25	29.41	25.59	24.34	12.86	13.48	27.24	19.43	15	17.28	15.21	15.63
∑PUFA	22.86	20.64	19.97	22.42	35.15	34.39	24.12	22.75	27.07	24.54	27.91	28.6
∑HUFA	4.24	4.79	4.68	4.01	4.72	6.87	5.54	3.67	6.21	6.89	6.97	12.44
∑(n-3)	16.97	14.92	12.1	14.6	16.09	17.74	19.12	9.83	18.82	18.24	21.19	24.81
∑(n-6)	9.08	9.18	10.85	10.77	22.50	21.92	10.49	10.49	14.31	11.07	13.56	15.97
(n-3/n-6)	1.86	1.62	1.11	1.35	0.71	0.8	1.82	0.93	1.31	1.64	1.56	1.55
DHA/EPA	0.04	0.03	0.02	0.02	0.02	0.3	0.02	0.1	0.01	0.03	6.83	0.01
Total lipid	8.41%	10.57%	8.24%	9.67%	4.91%	8.47%	5.79%	7.28%	7.23%	9.45%	9.41%	8.80%

Discussion

Variance analyses showed that cell density in light intensity 2500 and 6500 lux was significantly different ($P < 0.05$). Increasing light intensity increased cell density of *T. chuii*. With increasing light intensity, the photosynthetic rate also increased (Mendoza *et al.*, 1999). In fact, researchers believe that the light intensity until the chlorophyll II molecule is damaged can cause increased cell division (Thompson *et al.*, 1990; Bolch, 2004). This result agrees with those of Carvalho *et al.*, 2003. This micro alga in 6500 lux light intensity and 25 ppt salinity, most cell density induced, similar to result by Garcia *et al.*, 2007. That its optimum growth always occurs at salinity (25 ppt), on this regard, reduced growth rates at 40 ppt, because faster nutrient depletion in the culture medium (Johnson *et al.*, 1968). NaCl the essential element necessary for growth marine microalgae, thus over increase cause damage to cell microalgae. Rodolfi *et al.*, 2009 and Muller-Feuga *et al.*, 2003 also described in low salinity, increased total lipid content. Fatty acid composition depend to their cell growth (De la Pena *et al.*, 2005). Therefore, increase polyunsaturated fatty acid in high light intensity by reason of increase cell density. Our results on fatty acid showed that an increase in light intensity was associated with increased PUFA, HUFA, n-3, n-6 and decreased MUFA. On the other hand, Molina-Grima *et al.*, 1994 also described Decreases in the content of EPA with increasing light, in Batch cultures of *Isochrysis galbana*. It is widely accepted that low levels of light intensity bring about increases in the amount of thylakoid membranes, thus promoting synthesis of its Lipid constituents-galactolipids (which contain a high percentage of EPA). Therefore, any comparison of the intensity of light, they respond differently. Oleic acid, linoleic acid, alpha linolenic acid, an important fatty acids are unsaturated fatty acid (Pratoomyot *et al.*, 2005; Patil *et al.*, 2007). The significant presence of polyunsaturated fatty acids in these species, the nutritional value of food fish has doubled. Arachidonic acid and EPA are precursors of eicosanoid compounds. However, the eicosanoids from these two fatty acids are different both structurally and functionally, and are sometimes even antagonistic in their effects (Gill and Valivety, 1997). High ratios of n-3 to n-6 polyunsaturated fatty acids have been used as an index of high nutritional value to aquaculture animals (Watanabe *et al.*, 1983). The percentage differences in fatty acid composition in our study are similar to those reported by Pratoomyot *et al.*, 2005 for the *Tetraselmis sp* grown in laboratory culture with irradiances of 4719 lux. One difference was that they did not find DHA.

According to the result, a salinity of 25 ppt, light intensity 6500 lux seems more adequate for enhanced growth and fatty acid composition of *T. chuii*.

References:

- Bolch, C. 2004a. Lecture notes: KQA 201 - *Intensive Algal Culture*. School of Aquaculture, University of Tasmania, Launceston, Australia.
- Bold, H.C. and Wynne, M.J. 1985. Introduction to the algae structure and reproduction. Prentice-Hall Inc., Englewood Cliffs, New Jersey. 720 P.
- Brown, M.R., Dunstan, G.A., Norwood, S.J. and Miller, K.A. 1996. Effect of harvest stage on the biochemical composition of the diatom *Thalassiosira pseudonana*. *Journal of Phycology*, 32: 64-73.
- Carvalho, A.P. and Malcata, F.X. 2003. Kinetic modeling of the autotrophic growth of *Pavlova lutheri*: study of the combined influence of light and temperature. *Biotechnology Progress*, 19: 1128-1135.
- Creswell, L. 2010. Phytoplankton Culture for Aquaculture Feeds. *Proceedings of the 7th International Regional Aquaculture Center*, PP 1-12.
- De la Pena, M. and Villegas, C. 2005. Cell growth, effect of filamentous algae and nutritive value of the tropical Prasinophyte *Tetraselmis tetraele* at different phases of culture. Blackwell publishing Ltd, *aquaculture Research*, 36: 1500-1508.
- Desvilettes, C., Bourdier, G. and Breton, J.C. 1994. Lipid class and fatty acid composition of planktivorous larval pike (*Esox lucius*) living in a natural pond. *Aquatic Living Resources*, 7: 67-77.
- Fabregas, J., Abalde, J. and Herrero, C. 1984. Growth of the marine microalga *Tetraselmis suecica* in batch cultures with different salinities and nutrient concentrations. *Aquaculture*, 42: 207-215.
- Folch, J.M., Less, M. and Sloane-Stanley, G.H. 1957. A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biological Chemistry*, 226: 497-506.
- Garcia, F., Freile-Pelegrin, Y. and Robledo, D. 2007. Physiological characterization of *Dunaliella sp.* Chlorophyta, Volvocales from Yucatan, Mexico. *Bioresource Technology*, 98: 1359-1365

- Gill, I. and Valivety, R .1997. Polyunsaturated fatty acids: Part 1.Occurrence, biological activities and application. *Trends Biotechnology*,15:401–409.
- Johnson, M.K., Johnson, E.J., Mac Elroy, R.D., Speer, H.L. and Bruff, B.S. 1968. Effects of salts on the halophilic alga *Dunaliella viridis*. *Journal Bacteriology*, 95:1461–1468.
- Lartigue, J., Neill, A., Hayden, B.L., Pulfer, J. and Cebrian, J. 2003. The impact of salinity fluctuations on net oxygen production and inorganic nitrogen uptake by *Ulva lactuca* (Chlorophyceae). *Aquatic Botany*,75: 339-350.
- Makridis,P., Costa, R.A. and Dinis,M.T. 2006. Microbial conditions and antimicrobial activity in cultures of *Chlorella minutissima* and effect on bacterial load of enriched *Artemia metanauplii*. *Aquacult*, 255: 76-81.
- Mendoza, H., Martel, A., Jimenez, Del Rio M. and Garcia-Reina, G .1999. Oleic acid is the main fatty acid related with carotenogenesis in *Dunaliella salina*. *Journal Applied Phycology*, 11:15–19.
- Meseck, L.S.T., Jennifer, H.A. and Wikfors,H.G. 2005. Photoperiod and light intensity effects on growth and utilization of nutrients by the aquaculture feed microalgae, *Tetraselmis chuii*. *Aquaculture* 246, 393–404.
- Moisander, P. H., McClinton, E., and Paerl, H. W. 2002. Salinity effects on growth, photosynthetic parameters, and nitrogenase activity in estuarine planktonic cyanobacteria. In press, *Microbial Ecology*.
- Molina-Grima,E.,Camacho, F.G., Perez, J.A.S. and Sanchez-Gonzalez, J. 2004. Biochemical productivity and fatty acid profiles of *Isochrysis galbana* Parke and *Tetraselmis sp.* as a function of light intensity. *Process Biochemistry*, 29: 119–126.
- Muller-Feuga, A., Moal,J. and Kaas,R. 2003. The Microalgae of Aquaculture. J.G. Stottrup and L.A. McEvoy (eds.), *Live Feeds in Marine Aquaculture*, Blackwell Publishing, Oxford, PP.206-252.
- Patil,V., Kallqvist,T., Olsen,E., Vogt,G. and Gislerod,H.2007. Fatty acid composition of 12 microalgae for possible use in aquaculture feed. *Aquaculture International*, 15 :1-9.
- Pratoomyot,J., Srivilas,P. and Noiraksar,T.2005 . Fatty acids composition of 10 microalgal species. *Journal of Science and Technology*, 27: 1179-1187.
- Rodolfi, L., Zittelli, G. C., Bassi, N., Padovani, G., Biondi, N., Bonini, G. and Tredici, M. R. 2009. Microalgae for oil: Strain selection, induction of lipid synthesis and outdoor mass cultivation in a low-cost photobioreactor. *Biotechnology and Bioengineering* , 102:100-112.
- Seemann, J.R. and Critchley, C. 1985. Effect of salt stress on the growth, ion content, stomatal behavior and photosynthetic capacity of a salt-sensitive species, *Phaseolus vulgaris*. *Planta*, 164: 151-162.
- Smith, G.M. 1955. *Cryptogamic Botany, Volume I, Algae and Fungi*. McGraw-Hill Book Company, Inc. New York. 546P.
- Thompson, P. Harrison, P. and Whyte, J. 1990. Influence of irradiance on the fatty acid composition of phytoplankton. *Journal of Phycology* ,26: 278-288.
- Thompson, Jr., G.A. 1996. Lipids and membrane function in green algae. *Biochimica. Biophysica. Acta*, 1302:17– 45.
- Watanabe, T., Kitajima, C, and Jujita, S .1983. Nutritional value of live organisms used in Japan for mass propagation of fish: a review. *Aquaculture*: 34: 115-143.
- Zaki, M.L and Saad, H .2010. Comparative study on growth and survival of larval and juvenile *Dicentrarchus labrax* rearing on rotifer and *Artemia* enriched with four different microalgae species. *African Journal of Biotechnology*, 9: 3576-3588.